


Newly ice-free coastal zones as emerging carbon sinks in the warming Arctic fjords (Svalbard, West Spitsbergen)

Marlena Szeligowska
IO PAN

Polar Symposium BIS
19.05.2023
Sopot



Norway grants



GLACIERS RETREAT AND FIORDS WITHER

Gains and losses in the productivity of the polar fjords

euphotic zone ↓



*Fig. 1 Brown zones in front of
Tunabreen (14th Aug 2021)*



Norway
grants

Gains and losses in the productivity of the polar fjords

euphotic zone ↓

new area ↑

water mass exchange ↓




Fig. 3 Landsat RGB composites of Brepollen from 23rd July 1978 and 4th August 2022.

Gains and losses in the productivity of the polar fjords

euphotic zone ↓

new area ↑

water mass exchange ↓ ?

stratification ↓

nutrient balance ↓ ?

Fig. 4 The brown water caused by sediment being dredged up from the base of the glacier by meltwater plumes. Credit: NASA/JPL-Caltech



Gains and losses in the productivity of the polar fjords

euphotic zone ↓

new area ↑

water mass exchange ↓ ?

stratification ↓

nutrient balance ↓ ?

longer productive season ↑

advection ↑

Fig. 5 Arctic animals such as polar bears rely on sea ice that is shrinking as global temperatures rise. Credit: Ekaterina Anismova/AFP via Getty



Objective

Estimate the primary and zoobenthic production and carbon burial in the West Spitsbergen coastal waters and the newly ice-free area's contribution

Tab. 1 Primary production, zoobenthos and carbon burial in the West Spitsbergen fjords.

Variable	KGF outer	KGF inner	HOR outer	HOR inner	BIL	Reference
Summer pelagic primary production [mgCm ⁻² day ⁻¹]	108	59 80 - 155	336 - 1333	173		Piwoż et al., 2009 Iversen and Seuthe, 2011
Spring pelagic primary production [mgCm ⁻² day ⁻¹]		405 - 445 30 - 1850	320 - 2770		42.6	Iversen and Seuthe, 2011 Hodal et al., 2012 Vonnahme et al., 2021
Zoobenthos production [gCm ⁻² year ⁻¹]	9.4		19.2			Włodarska-Kowalczyk et al., 2019
Burial rate of OC [gCm ⁻² year ⁻¹]	28 ± 6		28 ± 1			Włodarska-Kowalczyk et al., 2019
	13	9				Kuliński et al., 2014
	10.0	5.7	19.3	30.3		Koziorowska et al., 2018
		15		38		Zaborska et al., 2018




Fig. 6 Map of the Svalbard archipelago with the investigated coastal zones.

Marine-terminating glaciers (1976-2022)

Data: Landsat satellite images

Time: summer 1976-2022

Resolution: 15 – 60 m

KKS – 15 glaciers
ISF – 8 glaciers

Fig. 7 Glacial bays with glaciers or glacial systems connected to the sea at least at one point in 1976 – 2022. Red dots represent SST data points. Pink dots represent the location of the meteorological stations. Background: Landsat8 satellite images from 27th July 2020



Marine-terminating glaciers (1976-2022)

Data: Landsat satellite images

Time: summer 1976-2022

Resolution: 15 – 60 m

VMK – 4 glaciers

HST – 15 glaciers

Fig. 8 Glacial bays with glaciers or glacial systems connected to the sea at least at one point in 1976 – 2022. Red dots represent SST data points. Pink dots represent the location of the meteorological stations. Background: Landsat8 satellite images from 4th August 2020





Norway
grants

Sea ice


Data: Landsat satellite images

Time: May 2022

Resolution: 15 m

KKS
ISF

Fig. 9 Sea ice in the West Spitsbergen fjords. Red dots represent SIC data points. Background: Landsat8 satellite images from May 2022





Norway
grants

Sea ice

Data: Landsat satellite images

Time: May 2022

Resolution: 15 m


VMK
HST

Fig. 10 Sea ice in the West Spitsbergen fjords. Red dots represent SIC data points. Background: Landsat8 satellite images from May 2022




Changes in the coastal zones area related to marine-terminating glaciers dynamics (1976-2022)

Fig. 11 Map of the Svalbard archipelago with the investigated coastal zones (left). The total area of the coastal zones, changes in the area in 1976-2022, and sea-ice cover in 2022 (right). Land and glaciers extent downloaded from <https://geodata.npolar.no/>.



Primary production, zoobenthic production and carbon burial

Fig. 12 Map of the Svalbard archipelago with the investigated coastal zones (left). Primary production (PP), zoobenthic production (ZBP), and organic carbon (OC) burial in the total area of the coastal zones (a) and newly ice-free areas (b).




Long-term trends in the coastal zones area, sea ice cover and melt potential

Data: Arctic Sea and Ice Surface Temperature, L4,
5km daily (DMI-ARC-SEAIC_TEMP-L4-NRT-OBS)
Time: 1976-2022
Resolution: 1 km, daily

Data: Meteorological stations (01003, 01007,
01008, Sveagruva)
Time: 1975-2022
Resolution: daily

Fig. 13 Long-term trends in the West Spitsbergen coastal waters: changes in the area related to marine-terminating glaciers dynamics, sea-ice duration, PDD SST and AT, melt season duration (* - $p < 0.05$, ** - $p < 0.001$ for modified Mann Kendall test).



Conclusions

more data needed

OC burial in the newly ice-free areas in the West Spitsbergen – only a small fraction of the global C burial in marine sediments

scale of marine ice loss worldwide

Fig. 14 The growing potential of Antarctic blue carbon (Sands et al., 2023)

